

COUNTEREXAMPLES FOR LOCAL ISOMETRIC EMBEDDING

NIKOLAI NADIRASHVILI AND YU YUAN

1. INTRODUCTION

In this paper, we construct metrics on 2-manifold which cannot be even locally isometrically embedded in the Euclidean space $\mathbb{R}^3$. By isometric embedding of (M^2, g) with $g = \sum_{i,j=1}^2 g_{ij} dx_i dx_j$ in $\mathbb{R}^3$, we mean there exists a surface in $\mathbb{R}^3$ with the induced metric equaling g , namely, the three coordinate functions $(X(x_1, x_2), Y(x_1, x_2), Z(x_1, x_2))$ defined on M^2 satisfy

$$dX^2 + dY^2 + dZ^2 = \sum_{i,j=1}^2 g_{ij} dx_i dx_j.$$

To be precise, we state the results in the following

Theorem 1.1. *There exists a smooth metric g in $B_1 \subset \mathbb{R}^2$ with Gaussian curvature $K_g \leq 0$ such that there is no C^3 isometric embedding of $(B_r(0), g)$ in $\mathbb{R}^3$ for any $r > 0$.*

Theorem 1.2. *There exists a smooth metric g in $B_1 \subset \mathbb{R}^2$ with Gaussian curvature $K_g(0) = 0$ and $K_g(x) < 0$ for $x \neq 0$ such that there is no $C^{3,\alpha}$ isometric embedding of $(B_r(0), g)$ in $\mathbb{R}^3$ for any $r > 0$ and $\alpha > 0$.*

Pogorelov [P2] constructed a simple $C^{2,1}$ metric g in $B_1 \subset \mathbb{R}^2$ with sign-changing Gaussian curvature such that (B_r, g) cannot be realized as a C^2 surface in $\mathbb{R}^3$ for any $r > 0$. Recently the first author [N] gave a C^∞ metric g on B_1 with no smooth isometric embedding of (B_r, g) in $\mathbb{R}^3$ for any $r > 0$. The sign of the Gaussian curvature K_g also changes.

On the positive side, when the sign of K_g for any smooth metric g does not change, the local smooth isometric embedding was settled by Pogorelov [P1], Nirenberg [Ni], and Hartman and Winter [HW2]. When $K_g \geq 0$ for the C^k metric with $k \geq 10$, there is a C^{k-6} isometric embedding of (B_{r_k}, g) in $\mathbb{R}^3$, this was done by Lin [L1]. When K_g changes sign cleanly, namely, $K_g(0) = 0, \nabla g(0) \neq 0$ for a C^k metric g , Lin [L2] showed that there exists a C^{k-3} isometric embedding in $\mathbb{R}^3$ for (B_{r_k}, g) with $k \geq 6$. When $K_g \leq 0$ and $\nabla^2 K_g(0) \neq 0$ for the smooth metric g , there is a local smooth isometric

Date: July 3, 2002.

Both authors are partially supported by NSF grants, and the second author also by a Sloan Research Fellowship.

embedding of g in $\mathbb{R}^3$, see Iwasaki [I]. When $K_g = -x_1^{2m} \tilde{K}(x)$ with $\tilde{K}(0) > 0$ for the smooth metric g , the same local isometric embedding also holds, see Hong [H]. Recently, Han, Hong, and Lin [HHL] showed that the local isometric embedding exists under the assumption $K_g \leq 0$ with a certain non-degeneracy of the gradient of K_g , or $K_g \leq 0$ with finite order vanishing.

If one allows higher dimensional ambient space, say $\mathbb{R}^4$, Poznyak [Po1] proved that any smooth metric g on M^2 can be locally smoothly isometrically embedded in $\mathbb{R}^4$. In fact, any C^k metric on n -manifold M^n has a C^k global isometric embedding in $\mathbb{R}^{N_n}$ with N_n large for $3 \leq k \leq \infty$. This is the work by Nash [Na2].

If we start with an analytic metric g on M^n , one always has a local analytic isometric embedding of (M^n, g) in $\mathbb{R}^{n(n+1)/2}$. This was proved by Janet [J], Cartan [C] very earlier on, and initiated by Schlaefli in 1873!

Lastly, any C^0 metric g on a compact n -manifold M^n which can be differentially embedded in $\mathbb{R}^{n+1}$ has a C^1 isometric embedding in $\mathbb{R}^{n+1}$, see Nash [Na1] and Kuiper [K].

For general description and further results on isometric embedding problem, we refer to [GR], [P2] and [Y].

The heuristic idea of the construction is to arrange the metric g in B_1 so that the second fundamental form of any isometric embedded surface in $\mathbb{R}^3$, $\text{II} \circ i$ vanishes at one point, where $i : (B_1, g) \rightarrow \mathbb{R}^3$ is the isometric embedding which is supposed to exist. Further we force $\text{II} \circ i$ to vanish along the boundary of a small domain Ω near the center of B_1 , where the Gaussian curvature $K_g < 0$ (in Ω). By the maximal principle, one cannot have a saddle surface with vanishing second fundamental form along the boundary. So (Ω, g) cannot be realized in $\mathbb{R}^3$. We repeat the construction near the center of B_1 at every scale so that (B_1, g) is not isometrically embeddable in $\mathbb{R}^3$ near the center.

The way to force $\text{II} \circ i$ to vanish at one point, say o , is the following. We modify the flat metric $g_0 = dx^2$ in $\mathbb{R}^2$ only over certain region Λ slightly away from the center o to a new one g so that, for a segment $A_1 A_2$ with $A_1, A_2 \in \partial\Lambda$, the length of $A_1 A_2$ under g is shorter than the one of the geodesic $A_1 A_2$ under the flat g_0 , and $K_g \leq 0$ in a subregion Λ_s containing $A_1 A_2$. Because of $\det \text{II}(i(0)) = 0$, we only need to deal with the other principle curvature. Suppose the second one $\kappa_2 \neq 0$, say $\kappa_2 < 0$. We show that there is a flat concave cylinder Σ near $i(B_1)$, which is isometric to (B_1, g_0) provided the embedding i is C^3 (This assertion for C^2 embedding case remains unclear to us). Now $i(A_1 A_2)$ supported on the saddle surface $i(\Lambda_s)$ can only stay above the concave cylinder Σ . Then the length of $i(A_1 A_2)$ is longer than the one of the projection of $i(A_1 A_2)$ down to the flat Σ , call it $P \circ i(A_1 A_2)$. We know the length of $P \circ i(A_1 A_2)$ under g_0 is equal to or longer than that of the geodesic $A_1 A_2$ under g_0 . But we start from $A_1 A_2$ with shorter length under g than under g_0 . This contradiction shows that $\text{II} \circ i(0)$ vanishes.

Inevitably, K_g is positive somewhere in Λ if Λ is surrounded by flat region with metric dx^2 . We add “tails” extending to the boundary ∂B_1 for the

modifying regions Λ , modify the metric on the tails, then we have the g with $K_g \leq 0$ in B_1 . It turns out that we cannot work with a segment in the construction, we go with a minimal tree connecting three points on $\partial\Lambda$ for each Λ , see section 2 for details.

Now that we have a non-isometrically embeddable metric (with nonpositive Gaussian curvature), the nearby metrics are almost non-isometrically embeddable. Based on this observation, we construct a non-isometrically embeddable metric with negative Gaussian curvature except for one point in section 3.

2. METRIC WITH NONPOSITIVE CURVATURE

Recall any three segments in $\mathbb{R}^2$ with equal angles $\frac{2}{3}\pi$ at the common vertex form a minimal tree T , namely, the length of T is less than that of any arcs connecting the other three vertices.

Lemma 2.1. *Let $u = -\operatorname{Im} e^{\log^2 z} = -e^{\log^2 r - \theta^2} \sin(2\theta \log r)$, $0 < \theta < 2\pi$. Then there exists a large integer K such that*

$$\int_T u ds < 0,$$

where the minimal tree $T = AA_1 \cup AA_2 \cup AA_3$ with $A = (-e^{-K}, 0)$, $A_2 = (-1, 0)$, $A_1, A_2 \in \partial B_1$, $\angle A_1AA_2 = \angle A_2AA_3 = \frac{2}{3}\pi$. Moreover, $u_r < 0$ for $r = 1$.

Proof. Set $\Omega_u = B_1 \cap \text{Sector } A_1AA_2$, $\Omega_l = B_1 \cap \text{Sector } A_2AA_3$, $\widehat{A_1A_2} = \partial\Omega_u \cap \partial B_1$, $\widehat{A_2A_3} = \partial\Omega_l \cap \partial B_1$. Let the angle from A_1A to x be φ , or $\varphi(x) = \angle A_1Ax$, then $0 \leq \varphi(x) \leq \frac{4}{3}\pi$ for $x \in \Omega_u \cup \Omega_l$.

We apply Green formula to harmonic functions u and φ in Ω_u and Ω_l ,

$$\begin{aligned} \int_{\partial\Omega_u} u \varphi_\gamma ds &= \int_{\partial\Omega_u} \varphi u_\gamma ds \\ \int_{\partial\Omega_l} u \left(\varphi - \frac{4}{3}\pi \right)_\gamma ds &= \int_{\partial\Omega_l} \left(\varphi - \frac{4}{3}\pi \right) u_\gamma ds, \end{aligned}$$

where γ is the outward unit normal of the integral domain. We then have

$$\begin{aligned} \int_{AA_1} -uds + \int_{AA_2} uds &= \int_{\widehat{A_1A_2}} \varphi u_r ds + \int_{AA_2} \frac{2}{3}\pi u_\theta ds \\ \int_{AA_2} -uds + \int_{AA_3} uds &= \int_{\widehat{A_2A_3}} \left(\varphi - \frac{4}{3}\pi \right) u_r ds + \int_{AA_2} \frac{2}{3}\pi u_\theta ds. \end{aligned}$$

It follows that

$$\begin{aligned} \int_{AA_1 \cup AA_3} uds &= 2 \int_{AA_2} uds + \int_{\widehat{A_1A_2}} -\varphi u_r ds + \int_{\widehat{A_2A_3}} \left(\varphi - \frac{4}{3}\pi \right) u_r ds \\ &= 2 \int_{AA_2} uds + \int_{\widehat{A_1A_2}} \varphi e^{-\theta^2} 2\theta ds + \int_{\widehat{A_2A_3}} \left(\frac{4}{3}\pi - \varphi \right) e^{-\theta^2} 2\theta ds. \end{aligned}$$

On the other hand,

$$\begin{aligned} \int_{AA_2} u ds &= \int_{e^{-K}}^{e^0} -e^{(\log^2 r - \pi^2)} \sin(2\pi \log r) dr \\ &= \frac{1}{2\pi e^{\pi^2}} \int_{-2\pi K}^0 -e^{\left(\frac{t^2}{4\pi^2} + \frac{t}{2\pi}\right)} \sin t dt. \end{aligned}$$

We choose large enough integer K so that $\int_{AA_2} u ds < 0$ and

$$2 \int_{AA_2} u ds + \int_{\widehat{A_1 A_2}} \varphi e^{-\theta^2} 2\theta ds + \int_{\widehat{A_2 A_3}} \left(\frac{4}{3}\pi - \varphi\right) e^{-\theta^2} 2\theta ds < 0.$$

Therefore

$$\int_T u ds < 0.$$

□

Remark. By applying Green formula to the above harmonic function u and linear functions, one sees that $\int_\Gamma u ds > 0$ for any segment $\Gamma \subset \Omega_u \cup \Omega_l$, connecting two boundary points on ∂B_1 .

Lemma 2.2. *There exists a function $v \in C_0^\infty(B_{1.1})$ satisfying*

$$\begin{aligned} v &= 0 & \text{in } \{(x_1, x_2) \mid x_1 < 0.9\} \setminus B_1 \\ \Delta v &\geq 0 & \text{in } B_1 \\ \int_T v ds &< 0 \end{aligned}$$

where the minimal tree $T = CC_1 \cup CA_2 \cup CC_3$ with $A_2 = (-1, 0)$, $C = (-\frac{1}{10}e^{-K} - 0.8, 0)$, $C_1, C_3 \in \partial B_1$ and $\angle C_1 CA_2 = \angle A_2 CC_3 = \frac{2}{3}\pi$. Moreover $T \subset \{(x_1, x_2) \mid x_1 < -0.1\}$.

Proof. Set $D = (-e^{-2K}, 0)$, $D_1, D_2 \in \partial B_1$ with $\angle D_1 DA_2 = \angle A_2 DD_3 = \frac{2}{3}\pi$, and $D_4 = (20, x_2(D_3))$, $D_5 = (20, x_2(D_1))$. Set $\Omega_p = \text{Pentagon } D_1 DD_3 D_4 D_5$. Let w satisfy

$$\begin{aligned} \Delta w &= 0 & \text{in } \Omega_p \\ w &= u & \text{on } D_1 D \cup D_3 D \\ w &= 0 & \text{on } D_1 D_5 \cup D_3 D_4 \\ w &= N & \text{on } D_4 D_5 \\ w &= u & \text{in } B_1 \setminus \text{Sector } D_1 DD_3, \end{aligned}$$

where u is the one in Lemma 2.1.

We choose large enough N so that $w_\gamma > u_\gamma$ on $D_1 D \cup D_3 D$ and $w_\gamma > 0$ on $D_1 D_5 \cup D_3 D_4$, where γ is the inward unit normal of $\partial\Omega_p$ this time. (If one insists, we can smooth off $\partial\Omega_p$.)




FIGURE 1. Minimal tree inside the half ball.

Next we mollify w by the usual (radially symmetric) mollifier $\rho_\delta \in C_0^\infty(B_\delta)$ with $0 < \delta < e^{-2K}$ to be determined later. We see that the smooth function $w * \rho_\delta$ satisfies

$$\Delta w * \rho_\delta(x) \geq 0 \quad \text{for } x_1 \leq 19.9$$

$$w * \rho_\delta(x) = u \quad \text{for } x \text{ inside } \Omega_i = B_1 \setminus \text{Sector } D_1 D D_3 \text{ and } \delta \text{ away from } \partial\Omega_i$$

$$w * \rho_\delta(x) = 0 \quad \text{for } x \text{ outside } \Omega_o = (B_1 \setminus \text{Sector } D_1 D D_3) \cup \Omega_p \text{ and } \delta \text{ away from } \partial\Omega_o.$$

Finally, set $C_0 = (-0.8, 0)$ and

$$v(x) = w * \rho_\delta(10(x - C_0)).$$

By making δ even smaller yet positive if necessary so that $\int_T v ds < 0$, we obtain the desired function v in the above lemma. $\square$

Corollary 2.1. *Let v be the function in Lemma 2.2. There exists a family of smooth metrics in $\mathbb{R}^2$*

$$g_\delta = e^{2\delta v} dx^2 \quad \text{for } 0 < \delta < \delta_0$$

such that

$$\begin{aligned} g_\delta &= dx^2 & \text{in } \{(x_1, x_2) \mid x_1 < 0.9\} \setminus B_1 \\ K_{g_\delta} &\leq 0 & \text{in } B_1 \\ L(T, g_\delta) &< L(T, dx^2), \end{aligned}$$

where $L(T, g)$ is the length of the minimal tree T from Lemma 2.2 in metric g .

Proof. We only prove the last two inequalities. One has

$$K_{g_\delta} = -e^{-2\delta v} \Delta(\delta v) \leq 0 \quad \text{in } B_1.$$

Also

$$L(T, g_\delta) = \int_T e^{\delta v} ds$$

$$\left. \frac{dL}{d\delta} \right|_{\delta=0} = \int_T v ds < 0.$$

Thus there exists δ_0 such that $L(T, g_\delta) < L(T, dx^2)$ for $0 < \delta < \delta_0$. $\square$

Let $\psi \in C^1([-1, 1])$ satisfy $0 \leq \psi \leq 1$ and $\psi(\pm 1) = 0$. Set $\gamma = \{(x_1, x_2) \mid x_1 = \psi(x_2), |x_2| \leq 1\}$, $Q = \{(x_1, x_2) \mid 0 < x_1 < \psi(x_2), |x_2| \leq 1\}$
 $\Pi = [0, 2] \times [-2, 2] \subset \mathbb{R}^2$, $F = \Pi \setminus Q$.

Lemma 2.3. *Let $f \in C^3(F)$. Assume the graph Σ of f is flat or $\det D^2 f = 0$ and $D^2 f \neq 0$ in F . Also assume a unit C^1 continuous eigenvector V_0 for the zero eigenvalue of $D^2 f$ is transversal to γ . For any $0 < \tau < 1$, there exists $\varepsilon > 0$ so that if $\left\| D^2 f - \begin{bmatrix} 0 & 0 \\ 0 & -\tau \end{bmatrix} \right\| \leq \varepsilon \tau$, one can extend f to Π with the graph of the extension being flat and concave.*

Proof. We take the C^2 Legendre coordinate system on $F \subset \Pi$ (cf. [HW1]).

$$\begin{cases} t = x_1 \\ s = f_2(x_1, x_2). \end{cases}$$

Notice that the graph of f , Σ is flat, or $\det D^2 f = 0$, it follows that $\{(x_1, x_2) \mid f_2(x_1, x_2) = s = \text{const}\}$ is a straight segment in $\mathbb{R}^2$ and $x_t(t, s)$ ($\|V_0\|$) is independent of t . Also $\frac{\partial f}{\partial t}(x(t, s))$ is independent of t . Hence we can represent a portion Σ^p of the graph Σ in the ruling form

$$(x_1, x_2, x_3)(t, s) = h(t, s) = c(s) + t\delta(s) = (t, x_2(t, s), f(t, x_2(t, s))),$$

where $c(s), \delta(s) \in C^2$ and $s \in S = [f_2(2, 2), f_2(2, -2)]$, $t \leq 2$.

We may assume $\nabla f(2, 0) = 0$. If ε is chosen small enough, then $\delta(s)$ ($\|V_0\|$) is close to $(1, 0, 0)$ in C^1 norm. Take ε small, then

$$\{(x_1, x_2, f(x_1, x_2)) \mid ((x_1, x_2) \in \gamma)\} \subset \partial \Sigma^p.$$

Set $U = \{(t, s) \mid -1 \leq t \leq 2, s \in S\}$. Take ε small so that $\|\delta(s) - (1, 0, 0)\|_{C^1}$ small, then $(t, s) \in U$ is a C^2 coordinate system for Π .

Now $\Sigma^e = h(U)$ is a C^2 , flat, concave graph over a domain Ω in $\mathbb{R}^2$ with $\Pi \subset \Omega$. Indeed, the normal of Σ^e is

$$N = \frac{h_t \times h_s}{\|h_t \times h_s\|}.$$

We know

$$h_t = \left(1, \frac{-f_{21}}{f_{22}}, f_1 + f_2 \frac{-f_{21}}{f_{22}}\right) \xrightarrow{\varepsilon \rightarrow 0} (1, 0, 0)$$

$$h_s = \left(0, \frac{1}{f_{22}}, \frac{f_2}{f_{22}}\right) \xrightarrow{\varepsilon \rightarrow 0} \left(0, \frac{-1}{\tau}, \frac{-s}{\tau}\right),$$

then $h_t \times h_s \xrightarrow{\varepsilon \rightarrow 0} (0, \frac{s}{\tau}, \frac{-1}{\tau})$. So Σ^e is a C^2 graph if we choose ε small enough.

Next, the second fundamental form of Σ^e is

$$\begin{aligned} II &= \begin{bmatrix} \langle h_{tt}, N \rangle & \langle h_{ts}, N \rangle \\ \langle h_{st}, N \rangle & \langle h_{ss}, N \rangle \end{bmatrix} \\ &= \frac{1}{\|h_t \times h_s\|} \begin{bmatrix} 0 & 0 \\ 0 & \langle c'' + t\delta'', \delta \times (c' + t\delta') \rangle \end{bmatrix} \end{aligned}$$

and the Gaussian curvature

$$K_g = 0.$$

Finally, the nonzero principle curvature of Σ^e

$$\kappa = \left[\frac{\tau^3}{(1+s^2)^{3/2}} + o(\varepsilon) \right] \langle c'' + t\delta'', \delta \times (c' + t\delta') \rangle.$$

On the other hand, from the graph representation of Σ^p , $\kappa \xrightarrow{\varepsilon \rightarrow 0} -\tau / (1+s^2)^{3/2}$. So for t in a certain range close to 2, say $t \in [1, 2]$, the quadratic function in terms of t ,

$$\langle c'' + t\delta'', \delta \times (c' + t\delta') \rangle = a_0 + a_1 t + a_2 t^2$$

is close to $-1/\tau^2$ as $\varepsilon \rightarrow 0$. It follows that $a_0 + a_1 t + a_2 t^2$ is still close to $-1/\tau^2$ for $t \in [-1, 2]$, if we choose ε small enough. So Σ^e is concave. $\square$

Lemma 2.4. *Let f be the extended function in Lemma 2.3, let $w \in C^2(\Pi)$ satisfy $w = f$ on F , $\det D^2 w \leq 0$ in Π , and $\left\| D^2 w - \begin{bmatrix} 0 & 0 \\ 0 & -\tau \end{bmatrix} \right\|_{C^1} \leq \varepsilon \tau$. Then*

$$f \leq w \text{ in } \Pi.$$

Proof. Suppose there is a point $x' = (x'_1, x'_2) \in M$ such that $w(x') < f(x')$. We know $x'_2 \in (-1, 1)$. For simplicity, we may assume

$$f(x') - w(x') = \sup_{x_2 \in [-1, 1]} [f(x'_1, x_2) - w(x'_1, x_2)].$$

Then $f_2(x') = w_2(x')$. It follows that the two tangent lines l_f, l_w to f and w at x' in the plane $\{(x_1, x_2, x_3) | x_1 = x'_1\}$ are parallel. Since $w(x'_1, \cdot)$ is concave, l_w is above w .

Let $T \subset \mathbb{R}^3$ be the tangent plane to the graph Σ_f of f at $(x', f(x'))$. Let $R = T \cap \Sigma_f$. Then R is a segment (ruling) transversal to l_f . Let $(x^0, z^0) \in R$ with $x^0 \in F$, then $z^0 = f(x^0) = w(x^0)$. Let $l_0 \subset T$ through (x^0, z^0) with $l_0 \parallel l_w$. By the concavity of $f = w$ in F , l_0 is above the graph Σ_w of w .

Let $m(x)$ be the linear function with graph as the plane E through l_w and l_0 . Let $V = \{(x_1, x_2) | x'_1 < x_1 < 2, |x_2| < 2\}$. Because Σ_w is a ruling surface on F , then

$$w(x) \leq m(x) \quad \text{on} \quad \partial V.$$

Note that $\det D^2w \leq 0$, by the maximum principle,

$$w(x) \leq m(x) \quad \text{in } V.$$

On the other hand, there is $(x^*, w(x^*)) \in R$ with $x^* \in V$ such that

$$w(x^*) > m(x^*).$$

This contradiction completes the proof of the above lemma. $\square$

Let r be a rotation in $\mathbb{R}^2$ through an angle 1° . Let v be the function in Lemma 2.2, set

$$w(x) = \sum_{i=1}^{360} v(r^i(1000x) - (360, 0)).$$

Pick two sequences $z_n \in \mathbb{R}^2$ and $\rho_n > 0$ such that

$$z_n \longrightarrow 0 \quad \text{as} \quad n \longrightarrow +\infty$$

$$B_{\rho_n}(z_n) \cap B_{\rho_k}(z_k) = \emptyset \quad \text{for} \quad n \neq k.$$

Take another sequence $\delta_n > 0$ going to 0 fast enough so that the smooth metric g_{II} in $\mathbb{R}^2$ satisfying

$$\begin{aligned} g_{II} &= e^{2\delta_n w(z_n + x/\rho_n)} dx^2 \quad \text{in } B_{\rho_n}(z_n) \\ g_{II} &= dx^2 \quad \text{otherwise.} \end{aligned}$$

Remark. Certainly our v is only smooth in $B_{1.1}(0)$, that leaves the function w nonsmooth, even undefined near the corresponding tails. At this stage, we do not need any information on the metric g_{II} near those tails (Figure 1 and 3). We can make a smooth extension of v to $\mathbb{R}^2$ with $v \in C_0^\infty(B_2)$ if one insists. Then the Gaussian curvature of g would be positive near the transition region. In the proof of Theorem 1.1, we will extend the tails to the boundary, make v a smooth subharmonic function inside the unit ball. Then the Gaussian curvature would be nonpositive in the unit ball.

Proposition 2.1. *Let i be a C^3 isometric embedding*

$$i : (B_r(0), g_{II}) \longrightarrow \mathbb{R}^3$$

for some $r > 0$. Then the second fundamental form of $i(B_r(0))$ vanishes at $i(0)$, or $II(i(0)) = 0$.

Proof. We may assume $i(B_r)$ is the graph Σ_w of a function $x_3 = w(x_1, x_2)$ and $w(0) = 0$, $\nabla w(0) = 0$. Then $II(i(0)) = D^2w(0)$ and $\det D^2w(0) = 0$. Suppose

$$D^2w(0) \neq 0.$$

Let P_3 be the projection from $\mathbb{R}^3$ to x_1 - x_2 plane. Set $J(x) = P_3(i(x))$. We may assume DJ is the identity map on the tangent space $\mathbb{R}^2$ at 0, and

$$D^2w(0, 0) = \begin{bmatrix} 0 & 0 \\ 0 & -\tau \end{bmatrix}.$$

For a sufficiently large n , $B_{\rho_n}(z_n) \subset B_r$ and

$$g_{II} = e^{2\delta_n v(r^{180}(1000(z_n + x/\rho_n)) - (360, 0))} dx^2$$

in the 179° to 181° section of the ball $B_{\rho_n}(z_n)$.

In order to simplify the presentation, we work with the metric $g_{\delta_n} = e^{2\delta_n v(x)} dx^2$ as in the Corollary 2.1. Let Σ^e be the flat, concave extension of $i(B_2^- \setminus B_1^-)$ by Lemma 2.3, where $B_\rho^- = \{(x_1, x_2) | x_1 < 0\} \cap B_\rho$. Note that we may consider the graph $x_3 = w_\varepsilon(x) = w(\varepsilon x)$ for small ε , then

$$\left\| D^2 w_\varepsilon - \begin{bmatrix} 0 & 0 \\ 0 & -\varepsilon^2 \tau \end{bmatrix} \right\|_{C^1} \leq \varepsilon^3,$$

make the extension, then scale back.

Since $i(B_1^-)$ is negatively curved, or $\det D^2 w \leq 0$ and concave, we apply Lemma 2.4 to conclude that $i(B_1^-)$ is above Σ^e .

Let P be the normal projection of points p above Σ^e down to Σ^e , that is $[p - P(p)] \perp \Sigma^e$. By concavity of Σ^e , we have

$$\text{Length}(T, g_{\delta_n}) = \text{Length}(i(T), g_{\Sigma_w}) \geq \text{Length}(P(i(T)), g_{\Sigma^e}),$$

Where g_{Σ_w} and g_{Σ^e} is the induced metrics on Σ_w and Σ^e .

Note that $P(i(C_1)) = i(C_1)$, $P(i(C_3)) = i(C_3)$, $P(i(A_2)) = i(A_2)$, there is an isometry $i_0 : \Sigma^e \rightarrow (\mathbb{R}^2, dx^2)$ such that $i_0 \circ P \circ i(C_1) = C_1$, $i_0 \circ P \circ i(C_3) = C_3$, $i_0 \circ P \circ i(A_2) = A_2$. Apply Corollary 2.1, we have

$$\text{Length}(P(i(T)), g_{\Sigma^e}) = \text{Length}(i_0 \circ P \circ i(T), dx^2) > \text{Length}(T, g_{\delta_n}).$$

Thus we arrive at

$$\text{Length}(T, g_{\delta_n}) > \text{Length}(T, g_{\delta_n}).$$

This contradiction finishes the proof of the above proposition. $\square$

Now we give the constructive proof of Theorem 1.1.

Proof. Step1. Let $\tilde{k}$ be a smooth function in $\mathbb{R}^2$ satisfying

$$\begin{aligned} \tilde{k} &< 0 \quad \text{in } B^n = B_{2^{-2n}}(2^{-n}, 0), \quad n = 1, 2, 3, \dots \\ \tilde{k} &= 0 \quad \text{otherwise.} \end{aligned}$$

Let u_1 be a smooth solution of

$$\Delta u_1 = -\tilde{k}.$$

Then the Gaussian curvature of the metric $g_1 = e^{2u_1} dx^2$ satisfies

$$\begin{aligned} K_{g_1} &= -e^{-2u_1} \Delta u_1 < 0 \quad \text{in } B^n \\ K_{g_1} &= 0 \quad \text{otherwise.} \end{aligned}$$

Step2. Choose a sequence $z_{n,k}$ outside each B^n and $\{(x_1, x_2) | x_2 = 0\}$ such that

$$\lim_{k \rightarrow \infty} z_{n,k} \in \partial B^n$$

$$\partial B^n \subset \overline{\{z_{n,k}\}_{k=1}^\infty}.$$

For each $z_{n,k}$, choose a simply connected thin tail $T_{n,k}$ with $T_{n,k}$ connecting $z_{n,k}$ and the boundary ∂B_1 such that

$$\begin{aligned} z_{n,k} &\in T_{n,k} \\ \partial T_{n,k} \cap \partial B_1 &= \text{a piece of arc with positive length} \\ T_{n,k} \subset \mathbb{R}_+^2 &= \{(x_1, x_2) | x_2 > 0\} \quad \text{for} \quad x_2(z_{n,k}) > 0 \\ T_{n,k} \subset \mathbb{R}_-^2 &= \{(x_1, x_2) | x_2 < 0\} \quad \text{for} \quad x_2(z_{n,k}) < 0 \\ T_{n,k} \cap T_{m,j} &= \emptyset \quad \text{for} \quad (n, k) \neq (m, j). \end{aligned}$$




FIGURE 2. Tails extending to the boundary.

We modify the metric $g_1 = e^{2u_1} dx^2$ over each tail $T_{n,k}$. But we proceed with the tails in the upper and lower half planes separately.

Since $K_{g_1} \equiv 0$ in the simply connected domain $\mathbb{R}_+^2 \setminus \cup_{n=1}^\infty B^n$. We represent $g_1 = dy_+^2$ in $\mathbb{R}_+^2 \setminus \cup_{n=1}^\infty B^n$ by a different coordinate system y_+ . Over each $T_{n,k} \subset \mathbb{R}_+^2$, we plant a metric

$$g_2 = e^{2V_{n,k}} dy_+^2 \quad \text{in} \quad x^{-1}(T_{n,k}),$$

where $V_{n,k}$ is similar to the one in the construction before Proposition 2.1, but the 360 disjoint sub-tails extend to the boundary $x^{-1}(\partial B_1)$ within




FIGURE 3. “Details” of one tail.

$x^{-1}(T_{n,k})$. We know $V_{n,k} = 0$ in $x^{-1}(B_1 \setminus T_{n,k})$. With $V_{n,k} = N_{n,k}$ chosen large enough on $x^{-1}(\partial B_1)$ intersection with the x pre-image of the 360 sub-tails, we make

$$\Delta V_{n,k} \geq 0 \quad \text{in} \quad x^{-1}(B_1).$$

We modify the metric $g_1 = e^{2u_1} dx^2$ over the tails in the lower half plane $\mathbb{R}_-^2$ with different coordinate system in the same way.

So far, we obtain a new metric $g_2 = e^{2u_2} dx^2$ in B_1 (which may not be smooth). We modify g_2 over the tails one last time.

Let

$$\begin{aligned} g_3 &= e^{2\epsilon_{n,k} V_{n,k}} dy_+^2 & \text{in} \quad x^{-1}(T_{n,k}) & \quad \text{for} \quad T_{n,k} \subset \mathbb{R}_+^2 \\ g_3 &= e^{2\epsilon_{n,k} V_{n,k}} dy_-^2 & \text{in} \quad x^{-1}(T_{n,k}) & \quad \text{for} \quad T_{n,k} \subset \mathbb{R}_-^2. \end{aligned}$$

By choosing $\epsilon_{n,k} > 0$, $\epsilon_{n,k} \rightarrow 0$ sufficiently fast for $k \rightarrow \infty$, we can assure $g_3 = e^{2u_3} dx^2$ is a smooth metric with $K_{g_3} \leq 0$ in B_1 .

Step 3. Suppose there is an isometric embedding

$$i : (B_r, g) \rightarrow \mathbb{R}^3$$

for some $r > 0$. Then there is n_* such that

$$B^{n_*} \subset B_r.$$

Applying Proposition 2.1, we have

$$II \circ i = 0 \quad \text{on} \quad \partial B^{n_*}.$$

We may assume $i(B_r)$ is represented as a graph $x_3 = f(x_1, x_2)$ with $\nabla f(0, 0) = 0$. Also we may assume the projection of $i(B^{n_*})$ down to x_1 - x_2 plane is a

domain Ω . Then

$$\begin{aligned} \det D^2 f &= K_g \left(1 + |\nabla f|^2\right)^2 < 0 \quad \text{in } \Omega \\ D^2 f &= 0 \quad \text{on } \partial\Omega. \end{aligned}$$

From $D^2 f = 0$ on $\partial\Omega$, it follows that $\nabla f = \text{const.}$ on $\partial\Omega$ and f coincides with a linear function on $\partial\Omega$. After subtracting the linear function from f , we may further assume $f = 0$ on $\partial\Omega$. We still have $\det D^2 f < 0$ in Ω . From the maximum principle, we see that $f \equiv 0$ in Ω . This contradiction finishes the proof of Theorem 1.1. $\square$

3. METRIC WITH NEGATIVE CURVATURE EXCEPT FOR ONE POINT

Relying on the metric constructed in Section 2, we construct a smooth metric g in B_1 with negative Gaussian curvature except for one point, namely, $K_g(x) < 0$ for $x \neq 0$, such that the surface (B_1, g) is not $C^{3,\alpha}$ isometrically embeddable in $\mathbb{R}^3$ even locally near 0.

For any surface (Ω, g) , we define the $C^{3,\alpha}$ isometric embedding norm by

$$\|(\Omega, g)\|_E = \inf \left\{ \|II(i(\Omega))\|_{C^{1,\alpha}} \mid C^{3,\alpha} \text{ isometric embedding } i : (\Omega, g) \longrightarrow \mathbb{R}^3 \right\}.$$

Now we give a constructive proof of Theorem 1.2.

Proof. Let the annulus $A^n = B_{1/n} \setminus B_{1/(n+1)} \subset \mathbb{R}^2$. We construct a metric $g = e^{2u_0} dx^2$ on B_1 such that a non-isometrically embeddable metric g as in Theorem 1.1 is planted (not just cut and pasted) over each annulus A^n .




FIGURE 4. Non-embeddable metric in each annulus.

Set

$$\widetilde{\varphi}_n(r) = \begin{cases} e^{-\frac{1}{r-1/n}} & r = |x| > \frac{1}{n} \\ 0 & 0 \leq r \leq \frac{1}{n} \end{cases}$$

We choose $\mu_1 > 0, \mu_2 > 0, \dots, \mu_n > 0, \dots$ such that $\varphi_n = \mu_n \widetilde{\varphi}_n$ satisfies that $\sum_{n=1}^{\infty} \varphi_n$ is smooth and even $\sum_{n=1}^{\infty} \epsilon_n \varphi_n$ is smooth for $(\epsilon_1, \epsilon_2, \dots) \in l^\infty$.

For $\epsilon = (\epsilon_1, \epsilon_2, \dots) \in l_+^\infty$, that is $\epsilon_1 > 0, \epsilon_2 > 0, \dots$ and $\|\epsilon\|_\infty = \max \epsilon_m < +\infty$, set

$$\Phi_\epsilon = \sum_{m=1}^{\infty} \epsilon_m \varphi_m$$

$$g_v = e^{2(u_0+v)} dx^2.$$

By the construction, $(A^n, e^{2u_0} dx^2)$ is not C^3 isometrically embeddable in $\mathbb{R}^3$ for any n , then we have the following.

There exists $0 < \eta_1$ such that $\|(A^1, g_{\Phi_\epsilon})\|_E \geq 1$ for $\epsilon \in l_+^\infty$ with $\|\epsilon\|_\infty \leq \eta_1$.

Next there exists $0 < \eta_2 < \eta_1$ such that $\|(A^m, g_{\Phi_\epsilon})\|_E \geq m$ for $m = 1, 2$ and $\epsilon = (\eta_1, \epsilon_2, \epsilon_3, \dots) \in l_+^\infty$ with $\|(0, \epsilon_2, \epsilon_3, \dots)\|_\infty \leq \eta_2$.

Inductively there exists $0 < \eta_k < \eta_{k-1}$ such that $\|(A^m, g_{\Phi_\epsilon})\|_E \geq m$ for $m = 1, 2, \dots, k$ and with $\epsilon = (\eta_1, \eta_2, \dots, \eta_k, \epsilon_{k+1}, \epsilon_{k+2}, \dots) \in l_+^\infty$ with $\|(0, \dots, 0, \epsilon_{k+1}, \epsilon_{k+2}, \dots)\|_\infty \leq \eta_k$.

...

Finally let $\Psi = \sum_{m=1}^{\infty} \eta_m \varphi_m$, $g = g_\Psi$. We see that

$$\|(A^m, g)\|_E \geq m \quad \text{for } m = 1, 2, 3, \dots$$

$$K_g(x) < 0 \quad \text{for } x \neq 0 \quad \text{and}$$

$$K_g(0) = 0.$$

It follows that there is no $C^{3,\alpha}$ isometric embedding of $(B_r(0), g)$ in $\mathbb{R}^3$ for any $r > 0, \alpha > 0$. $\square$

REFERENCES

- [C] Cartan, E., *Sur la possibilité de plonger un espace riemannien donné dans un espace euclidien*, Ann. Soc. Pol. Math., **5** (1927), 1–7.
- [GR] Gromov, M. L. and Rokhlin, V. A., *Embeddings and immersions in Riemannian geometry*, Uspehi Mat. Nauk, **25** (1970) 3–62; English translation in Russian Math. Survey **25** (1970), 1–57.
- [HHL] Han, Q., Hong, J. X., and Lin, C. S., to appear.
- [HW1] Hartman, P. and Winter, P., *On the asymptotic curves of a surface*, Amer. J. Math., **73** (1951), 149–172.
- [HW2] Hartman, P. and Winter, P., *On hyperbolic partial differential equations*, Amer. J. Math., **74** (1952), 834–864.
- [I] Iwasaki, N., *Applications of Nash-Moser theory to nonlinear Cauchy problems*, Proc. Sympos. Pure Math., **45** (1986) Part 1, 525–528.
- [H] Hong, J. X., *Cauchy problems for degenerate hyperbolic Monge-Ampère equations and some applications*, J. Partial Differential Equations **4** (1991), 1–18.
- [J] Janet, M., *Sur la possibilité de plonger un espace riemannien donné dans un espace euclidien*, Ann. Soc. Pol. Math., **5** (1926), 38–43.

- [K] Kuiper, N. H., *On C^1 -isometric embeddings, I, II*, Indag. Math., **17** (1955), 545–556, 683–689.
- [L1] Lin, C. S., *The local isometric embedding in $\mathbb{R}^3$ of 2-dimensional Riemannian manifolds with nonnegative curvature*, J. Diff. Geom., **21** (1985), 213–230.
- [L2] Lin, C. S., *The local isometric embedding in $\mathbb{R}^3$ of two-dimensional Riemannian manifolds with Gaussian Curvature changing sign cleanly*, Comm. Pure Appl. Math., **39** (1986), 867–887.
- [N] Nadirashvili, N., *The local embedding problem for surfaces*, preprint, 2002.
- [Na1] Nash, J., *C^1 isometric imbeddings*, Ann. of Math., **60** (1954), 383–396.
- [Na2] Nash, J., *The imbedding problem for Riemannian manifolds*, Ann. of Math., **63** (1956), 20–63.
- [Ni] Nirenberg, L., *The Weyl and Minkowski problems in differential geometry in the large*, Comm. Pure Appl. Math., **6** (1953), 337–394.
- [P1] Pogorelov, A. V., *Intrinsic estimates for the derivatives of the radius vector of a point on regular convex surface*, Dokl. Akad. Nauk SSSR (N.S.), **66** (1949), 805–808.
- [P2] Pogorelov, A. V., *An example of a two-dimensional Riemannian metric that does not admit a local realization in E^3* , Dokl. Akad. Nauk SSSR, **198** (1971) 42–43; English translation in Soviet Math. Dokl. **12** (1971), 729–730.
- [Po] Poznyak, È. G., *Isometric embeddings of two-dimensional Riemannian metrics in Euclidean space*, Uspekhi Mat. Nauk, **28** (1973), 47–76; English translation in Russian Math. Survey, **28** (1973), 47–77.
- [Y] Yau, S.-Y., *Seminar on Differential Geometry*, Annals of Math. Studies, **102**, Princeton, 1982.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, 5734 S. UNIVERSITY AVE., CHICAGO, IL 60637

E-mail address: nicholas@math.uchicago.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, 5734 S. UNIVERSITY AVE., CHICAGO, IL 60637, AND UNIVERSITY OF WASHINGTON, SEATTLE, WA 98195

E-mail address: yuan@math.uchicago.edu, yuan@math.washington.edu